



Chemotaxis Assays

Analyze Directed Migration of Adherent and Non-Adherent Cells

✓ Chemotaxis Measurement in Real-Time

Complete solution from sample preparation to quantitative data analysis

✓ Long-Term Experiments

Keep a linear and stable chemotactic gradient for over 48 hours

✓ Reproducible Results

Reliable and user-independent data

2D Chemotaxis

Chemotaxis of slow migrating, adherent cells on 2D surfaces (e.g., cancer cells, endothelial cells, and fibroblasts)

3D Chemotaxis

Chemotaxis of fast or slow migrating cells in 3D gel matrices (e.g., immune cells, tumor cells, and endothelial cells)


Additional equipment for researchers doing chemotaxis assays:








www.ibidi.com




Chemotaxis Assays

Analyze Directed Migration of Adherent and Non-Adherent Cells


Sample Preparation




Live Cell Imaging



Cell Tracking




Data Analysis



The Principle of the µ-Slide Chemotaxis

Using the μ -Slide Chemotaxis, a precisely defined and stable chemotactic gradient is created in a reproducible environment.



Live Cell Microscopy and Cell Tracking

With live cell imaging, chemotaxis can be analyzed under physiological conditions in real time. Tracking the cells allows quantification of their movement over time. It can be done manually or automatically with special tracking software.

Data Analysis

For further analysis of tracking data, ibidi has developed the **Chemotaxis and Migration Tool**, which is available for free at ibidi.com/chemotaxis-tool. This software provides statistical analysis and graphical display of the experimental data.



Register now for a **2-day laboratory course** at ibidi Munich, Germany: ibidi.com/practical-courses.

FREE SAMPLES: ibidi.com/free-samples

Technical Details:

| Chemotaxis chambers on slide | 3 |
|------------------------------|-----------------------|
| Volume per chamber | 120 μΙ |
| Observation area | 2 x 1 mm ² |
| Total height with plugs | 12 mm |
| Volume chemoattractant | 30 μΙ |

Ordering Information:

| Cat. No. | Description | Pcs./Box |
|----------|--|----------|
| 80326 | μ-Slide Chemotaxis ibiTreat: #1.5 polymer coverslip, tissue culture treated, sterilized | 10 |
| 80322 | μ-Slide Chemotaxis Collagen IV: #1.5 polymer coverslip, sterilized | 10 |
| 00000 | | 10 |
| 80328 | sticky-Slide Chemotaxis: sterilized | 10 |
| 10812 | Coverslips for sticky-Slides: #1.5H (170 µm +/- 5 µm) D 263 M Schott glass, 25 mm x 75 mm, unsterile | 100 |

ibidi GmbH Am Klopferspitz 19 82152 Martinsried Germany Tel.: +49 89 / 520 46 17 - 0 Fax: +49 89 / 520 46 17 - 59 E-Mail: info@ibidi.de

E-Mail: info@ibidi.de © ibidi GmbH, V5.0, 2018/02 For more information on our products, please visit us at:

